Nitrogen Oxides (NOx / NO2)

Regulatory History — CASRN 10102-44-0

Nitrogen Oxides (NOx)

Nitrogen oxides (NOx) are a group of highly reactive gases composed primarily of nitric oxide (NO) and nitrogen dioxide (NO2) that are produced during high-temperature combustion. These compounds are among the most common air pollutants emitted from both natural and anthropogenic sources. Major emission sources include motor vehicles, power plants, industrial boilers, petroleum refineries, chemical manufacturing facilities, internal combustion engines, marine vessels, aircraft, wildfires, prescribed burns, and structural fires. Nitric oxide is typically emitted directly from combustion processes and rapidly oxidizes in the atmosphere to form nitrogen dioxide, which subsequently participates in complex atmospheric reactions that produce ozone, nitrate particulate matter, and acid deposition. Because nitrogen oxides are chemically reactive and widely emitted, they play a central role in regional air pollution and atmospheric chemistry.

Scientific concern regarding nitrogen oxides increased throughout the twentieth century as urban air pollution became more prevalent and epidemiological studies demonstrated associations between NO2 exposure and adverse respiratory health effects. Research conducted during the development of photochemical smog control programs identified nitrogen oxides as critical precursors in the formation of ground-level ozone and fine particulate matter (PM2.5), two pollutants responsible for substantial public health impacts. Continued toxicological, clinical, and epidemiological investigations have established that nitrogen oxides contribute to respiratory disease, reduced lung function, and increased cardiovascular morbidity, making them among the most extensively studied criteria air pollutants.

Nitrogen dioxide (NO2) is regulated as one of the six criteria air pollutants under the Clean Air Act. The U.S. Environmental Protection Agency (EPA) has established National Ambient Air Quality Standards (NAAQS) that include an annual primary standard of 53 parts per billion (ppb) and a 1-hour primary standard of 100 ppb to protect public health. Nitrogen oxide emissions are also regulated through numerous federal programs, including New Source Performance Standards (NSPS), National Emission Standards for Hazardous Air Pollutants (NESHAP), vehicle emission standards, state implementation plans (SIPs), and ozone transport regulations. Occupational exposure is regulated by the Occupational Safety and Health Administration (OSHA), which has established a permissible exposure limit (PEL) of 25 ppm for

nitric oxide (NO) and 5 ppm as a ceiling limit for nitrogen dioxide (NO_2). Over recent decades, implementation of catalytic converters, low-NOx burners, selective catalytic reduction (SCR), and other emission control technologies has substantially reduced nitrogen oxide emissions from many major sources.

Nitrogen oxides have been extensively evaluated by regulatory and public health agencies because of their respiratory toxicity. The International Agency for Research on Cancer (IARC) has not classified nitrogen oxides as human carcinogens, and the National Toxicology Program (NTP) does not list nitric oxide or nitrogen dioxide as known or reasonably anticipated human carcinogens. Likewise, nitrogen oxides are not listed under California Proposition 65as chemicals known to cause cancer. Regulatory concern instead focuses on their ability to produce acute respiratory effects, contribute to chronic respiratory disease, and indirectly affect human health through the formation of ozone, nitrate aerosols, and secondary fine particulate matter.

Scientific understanding of nitrogen oxide toxicity has expanded considerably through toxicological, epidemiological, and controlled human exposure studies. Acute exposure to elevated nitrogen dioxide concentrations may produce coughing, wheezing, chest tightness, airway inflammation, shortness of breath, and reduced pulmonary function, particularly among individuals with asthma or other preexisting respiratory conditions. Repeated or long-term exposure has been associated with increased asthma exacerbations, respiratory infections, chronic bronchitis, impaired lung development in children, and elevated hospital admissions for respiratory and cardiovascular disease. In addition to their direct respiratory effects, nitrogen oxides contribute significantly to secondary pollutant formation, including ozone and $PM_{2.5}$, thereby extending their health impacts well beyond the initial emission source.

Nitrogen oxides remain among the most important contaminants evaluated in environmental forensic investigations because they are reliable indicators of combustion processes and major contributors to regional air pollution. Environmental investigations commonly evaluate nitrogen oxide emissions from power plants, petroleum refineries, industrial combustion equipment, transportation sources, marine vessels, airports, wildfires, prescribed burns, and structural fires. Forensic investigations often integrate continuous emissions monitoring, ambient air monitoring, satellite observations, atmospheric chemistry, emissions inventories, and air dispersion modeling to identify emission sources, quantify downwind concentrations, evaluate secondary pollutant formation, and assess community exposure. Because nitrogen oxides participate in complex atmospheric reactions that generate ozone and nitrate particulate matter, environmental investigations frequently consider both primary nitrogen oxide emissions and their secondary transformation products when evaluating air quality impacts and potential human health risks.

References

Agency for Toxic Substances and Disease Registry. (2018). Toxicological profile for nitrogen oxides (interaction profile). U.S. Department of Health and Human Services. https://www.atsdr.cdc.gov

International Agency for Research on Cancer. (2016). Outdoor air pollution (IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 109). World Health Organization. https://monographs.iarc.who.int/

National Toxicology Program. (2021). Report on carcinogens (15th ed.). U.S. Department of Health and Human Services. https://ntp.niehs.nih.gov/whatwestudy/assessments/cancer/roc

Occupational Safety and Health Administration. (n.d.). Nitric oxide and nitrogen dioxide.https://www.osha.gov/chemicaldata

U.S. Environmental Protection Agency. (2024). Nitrogen dioxide (NO_2) pollution. https://www.epa.gov/no2-pollution

U.S. Environmental Protection Agency. (2024). National Ambient Air Quality Standards (NAAQS) for nitrogen dioxide.https://www.epa.gov/no2-pollution/primary-national-ambient-air-quality-standards-naaqs-nitrogen-dioxide

U.S. Environmental Protection Agency. (2023). Integrated Science Assessment (ISA) for Oxides of Nitrogen – Health Criteria. https://www.epa.gov/isa

World Health Organization. (2021). WHO global air quality guidelines: Particulate matter ($PM_{2.5}$ and PM_{10}), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. World Health Organization. https://www.who.int/publications/i/item/9789240034228